

CONTINUOUS TIME MARKOV PROCESSES ON GRAPHS

JIANJUN TIAN AND XIAO-SONG LIN

ABSTRACT. We study continuous time Markov processes on graphs. The notion of frequency is introduced, which serves well as a scaling factor between any Markov time of a continuous time Markov process and that of its jump chain. As an application, we study “multi-person simple random walks” on a graph G with n vertices. There are n persons distributed randomly at the vertices of G . In each step of this discrete time Markov process, we randomly pick up a person and move it to a random adjacent vertex. We give estimate on the expected number of steps for these n persons to meet all together at a specific vertex, given that they are at different vertices at the begininng. For regular graphs, our estimate is exact.

1. INTRODUCTION

In this paper, for simplicity, we will consider connected simple graphs only. These are connected graphs without multiple edges and loops. We will adopt the following notations and terminologies for a graph G . The sets of vertices and edges of G are $V(G)$ and $E(G)$, respectively. The *order* n of G is the number of vertices of G , and the *size* m of G is the number of edges of G . Thus, $n = |V(G)|$ and $m = |E(G)|$. For a vertex $x \in V(G)$, $\Gamma(x)$ is the set of vertices which are connected to x by an edge in $E(G)$. The *degree* of a vertex x is $d(x) = |\Gamma(x)|$. We have

$$\sum_{x \in V(G)} d(x) = 2m.$$

The adjacent matrix of G is denote by $A(G)$ and the diagonal matrix $D(G)$ has the sequence of degrees at each vertex as its diagonal entries. Finally, we denote

$$d_m = \min \{d(x) ; x \in V(G)\} \quad \text{and} \quad d_M = \max \{d(x) ; x \in V(G)\}.$$

What concerns us primarily in this paper is the following continuous time Markov process on a graph G : The probability that a person standing at a vertex x of G to jump to a neighboring vertex in $\Gamma(x)$ within a time period Δt is $d(x)\Delta t + o(\Delta t)$, and once jumped, the person at x has equal probability to land at a vertex $y \in \Gamma(x)$. If we write

$$Q = Q(G) = -D(G) + A(G),$$

the transition probability matrix of this Markov process is

$$P(t) = e^{tQ}.$$

We call this Markov process CTSRW (continuous time simple random walks). In the literature, it is the discrete time simple random walks (SRW) on a graph G that people concern most. One usually call SRW the *jump chain* of CTSRW. The transition probability matrix of SRW is $D(G)^{-1}A(G)$.

We introduce in this paper a fundamental quantity for CTSRW on a graph G called *frequency*. Let $N(t)$ be the expected number of jumps of the Markov process CTSRW up to time t . Then the frequency f of CTSRW is defined to be

$$f = \lim_{t \rightarrow \infty} \frac{N(t)}{t}.$$

Using the Lévy formula, we are able to calculate the frequency for CTSRW and get

$$f = \frac{2m}{n}.$$

The frequency turns out to be a natural scaling factor between various important quantities of CTSRW and SRW, respectively. For example, we have the following theorem.

Theorem 1.1. *Let G be non-bipartite. For a vertex x of G , let T_x be the first return time of CTSRW on G and N_{T_x} be the number of jumps during the time period $[0, T_x]$. Notice that N_{T_x} is the first return time for the discrete time SRW. Then, the expectations of T_x and N_{T_x} , $E(T_x)$ and $E(N_{T_x})$, respectively, are related by the following equation:*

$$E(N_{T_x}) = fE(T_x).$$

More generally, we have the following theorem.

Theorem 1.2. *Let T be any Markov time (or stopping time) of CTSRW on a graph G with finite expectation, let N_T be the number of jumps during the time period $[0, T]$. Then*

$$d_m \leq f \leq d_M$$

and

$$d_m E(T) \leq E(N_T) \leq d_M E(T).$$

In particular, if G is regular so that $d_m = d_M$, we have $E(N_T) = fE(T)$.

As an application, we consider *multi-person simple random walks* (MPSRW) on a graph G . To start with, we assume that each of the n vertices of G is occupied by a person. In each step of this Markov chain, there is one person, equally possible for each of these n persons, who moves to a neighboring vertex, also equally possible for each of the neighboring vertices. We would like to know the expected number of steps this Markov chain should take for these n persons to meet all together at a specified vertex.

We will see that this Markov chain is the jump chain of a continuous time Markov process on the set M_n of *maps* from $\{1, 2, \dots, n\}$ to itself. On the other

hand, this continuous time Markov process on M_n can be identified with the n -th tensor power of CTSRW on G . Thus, computation of expectations of various stopping times for this continuous time Markov process on M_n can be carried out. We are then able to use Theorem 1.2 mentioned above to obtain estimates for the expected number of steps for MPSRW on G to coalesce.

We refer the reader to [2, 5, 6] for basic terminologies and results in the study of simple random walks on graphs.

2. CONTINUOUS TIME MARKOV PROCESS ON WEIGHTED GRAPHS

Let G be a connected weighted graph with order n size m , we define continuous time Markov process X_t on G by giving its infinitesimal generator Q as the negative weighted Laplacian of G . Specifically, every edge $xy \in E(G)$ is associated with a positive number (weight) w_{xy} . We do not direct edges in G and therefore $w_{xy} = w_{yx}$. We denote by

$$w_x = \sum_{y \in \Gamma(x)} w_{xy}$$

the total weight at the vertex x . We write $Q = (q_{xy})_{n \times n}$, where

$$q_{xy} = \begin{cases} w_{xy} & \text{if } xy \in E(G), \\ -w_x & \text{if } x = y, \\ 0 & \text{otherwise.} \end{cases}$$

Thus the probability transition matrix X_t is given by

$$P(t) = e^{tQ} = (p_{xy}(t))_{n \times n},$$

and transition probability from vertex x to vertex y is given by

$$\Pr \{X(h+t) = y \mid X(h) = x\} = p_{xy}(t).$$

By the way, we may call $-Q = L_w$ the weighted Laplacian of the weighted graph G .

In the special case of $w_{xy} = 1$ for all $xy \in E(G)$, we have CTSRW on the graph G . The infinitesimal generator $Q = -L(G) = -D(G) + A(G) = (q_{xy})_{n \times n}$ is given by

$$q_{xy} = \begin{cases} 1 & \text{if } xy \in E(G), \\ -d(x) & \text{if } x = y, \\ 0 & \text{otherwise.} \end{cases}$$

2.1. Ergodicity. We set $u = (\frac{1}{n}, \frac{1}{n}, \dots, \frac{1}{n})$ to be a probability vector. Then, since Q is symmetric, u is an invariant measure of the Markov process X_t . Namely, we have $uQ = 0$ and

$$uP(t) = u \left(I + tQ + \frac{t^2}{2!}Q^2 + \dots \right) = u.$$

We claim that u also is the ergodic vector or the stationary distribution. To see this, notice first that the graph G is connected so the process X_t is irreducible. Thus

$$\lim_{t \rightarrow \infty} p_{xy}(t) = v_y$$

exists and does not depend on x . Actually, we have $v_y = u_y = \frac{1}{n}$ by the following calculation: First, for any fixed $t > 0$,

$$\begin{aligned} uP(2t) &= uP(t)P(t) = uP(t) = u, \\ &\dots\dots\dots \\ uP(kt) &= uP((k-1)t) = \dots = uP(t) = u. \end{aligned}$$

Then

$$u_y = \sum_x u_x p_{xy}(kt), \quad \text{for } y \in V(G).$$

Let $k \rightarrow \infty$, we get

$$u_y = \sum_x u_x v_y = v_y.$$

2.2. The mean first return time. For a vertex $x \in V(G)$, denote by T_{xx} the first return time to x , given that the Markov process X_t starts at x . That is

$$T_{xx} = \inf \{t : t > \rho_x, X_t = x \mid X_0 = x\}$$

where ρ_x is the exit time from the vertex x . We denote by $h(x, x)$ the mean first return time $E(T_{xx})$.

Lemma 2.1. *The mean first return time is*

$$h(x, x) = \frac{n}{w_x}.$$

Proof. Write

$$F_{xx}(t) = \Pr \{T_{xx} \leq t\}.$$

Then we have the equation [4]

$$p_{xx} = e^{-w_x t} + \int_0^t p_{xx}(t-s) dF_{xx}(s).$$

Taking Laplace transform, we get

$$\begin{aligned}
\phi_{xx}(\lambda) &= \int_0^\infty e^{-\lambda t} p_{xx}(t) dt \\
&= \int_0^\infty e^{-(\lambda+w_x)t} dt + \int_0^\infty e^{-\lambda t} \int_0^t p_{xx}(t-s) dF_{xx}(s) dt \\
&= \frac{1}{\lambda + w_x} + \int_0^\infty \int_s^\infty e^{-\lambda t} p_{xx}(t-s) dt dF_{xx}(s) \\
&= \frac{1}{\lambda + w_x} + \int_0^\infty \int_0^\infty e^{-\lambda(v+s)} p_{xx}(v) dv dF_{xx}(s) \\
&= \frac{1}{\lambda + w_x} + \int_0^\infty e^{-\lambda s} \int_0^\infty e^{-\lambda v} p_{xx}(v) dv dF_{xx}(s) \\
&= \frac{1}{\lambda + w_x} + \int_0^\infty e^{-\lambda s} \phi_{xx}(\lambda) dF_{xx}(s) \\
&= \frac{1}{\lambda + w_x} + \phi_{xx}(\lambda) l_x(\lambda)
\end{aligned}$$

where $l_x(\lambda) = \int_0^\infty e^{-\lambda s} dF_{xx}(s)$. Then, we have

$$\lambda \phi_{xx}(\lambda) = \frac{1}{\lambda + d(x)} \left(\frac{1 - l_x(\lambda)}{\lambda} \right)^{-1}.$$

Since

$$\lim_{\lambda \rightarrow 0} \frac{1 - l_x(\lambda)}{\lambda} = \lim_{\lambda \rightarrow 0} \int_0^\infty s e^{-\lambda s} dF_{xx}(s) = \lim_{\lambda \rightarrow 0} \int_0^\infty s dF_{xx}(s) = h(x, x)$$

and

$$\lim_{\lambda \rightarrow 0} \lambda \phi_{xx}(\lambda) = u_x,$$

we get

$$\begin{aligned}
u_x &= \lim_{\lambda \rightarrow 0} \frac{1}{\lambda + w_x} \left(\frac{1 - l_x(\lambda)}{\lambda} \right)^{-1} \\
&= \frac{1}{h(x, x) d(x)}.
\end{aligned}$$

Thus,

$$h(x, x) = \frac{n}{w_x}.$$

□

2.3. The mean hitting time of y from x . In general, define

$$T_{xy} = \inf \{t : t > \rho_x, X_t = y \mid X_0 = x\}.$$

I. e. T_{xy} is the time of first entrance into, or hitting, the vertex y , given that the process starts at x . We denote the mean first hitting time of y from x by $h(x, y) = E(T_{xy})$. We have the following equation:

$$h(x, x) = \frac{1}{w_x} + \sum_{y \in \Gamma(x)} \frac{w_{xy}}{w_x} h(y, x) = \frac{1}{w_x} + \frac{1}{w_x} \sum_{y \in \Gamma(x)} w_{xy} h(y, x).$$

Lemma 2.2. *The mean hitting time can be calculated from the following integral:*

$$h(x, y) = n \int_0^\infty (p_{yy}(t) - p_{xy}(t)) dt.$$

The formula in this lemma is similar to the formula in the discrete time case. We omit the proof since it is also analogous to the discrete time case.

2.4. The stationary distribution of SRW. For an unweighted graph G , the jump chain CTSRW on G is SRW on G . We know that $Q = -D + A$, and the transition probability matrix of SRW is $D^{-1}A$. If we set $\tilde{\pi} = uD$, where u is the stationary distribution of CTSRW, then $uQ = -uD + uA = 0$. So, substitute $u = \tilde{\pi}D^{-1}$ we have

$$\tilde{\pi} = \tilde{\pi}D^{-1}A.$$

Thus, $\tilde{\pi}$ is an invariant measure of SRW. We need to normalize it. Let

$$\pi = \frac{1}{\sum_{i=1}^n \tilde{\pi}_i} \tilde{\pi}.$$

Then π is an invariant distribution for SRW. Specifically

$$\pi_x = \frac{d(x)}{\sum_{x \in V(G)} d(x)} = \frac{d(x)}{2m}.$$

If the graph G is non-bipartite, this invariant distribution is also the stationary distribution. It is well known that the mean number of steps SRW should take to return to the vertex x for the first time is $1/\pi_x = 2m/d(x)$. Recall that the mean first return time of CTSRW is $n/d(x)$. Therefore it is natural to think of the quantity

$$f = \frac{2m/d(x)}{n/d(x)} = \frac{2m}{n}$$

as the *frequency* (number of jumps per unit time) of CTSRW. We will make this notion precise in the following subsection.

2.5. The frequency. To define the frequency for the continuous time Markov process X_t on a weighted graph, we first define a quantity $N(t)$ for $t > 0$:

$$N(t) = E(\text{the number of jumps of } X_t \text{ up to time } t).$$

Theorem 2.1. *We have*

$$f := \lim_{t \rightarrow \infty} \frac{N(t)}{t} = \frac{2w}{n}$$

where w is the total weight of G

$$w = \frac{1}{2} \sum_{xy \in E(G)} w_{xy}.$$

Proof. Let us recall the Lévy formula first. See [4]. Given a Markov process X_t , we consider a purely discontinuous functional $A = \{A_t : 0 < t < \infty\}$ on the path space defined by

$$A_t = \sum_{0 < s \leq t} g(X_{s-}, X_s), \quad t > 0,$$

where g is a function on $V(G) \times V(G)$. Also, we define a function b_Q on $V(G)$ by

$$b_Q(x) = \sum_{y \neq x} w_{xy} g(x, y), \quad x \in V(G),$$

and the integral functional $B = \{B_t : 0 \leq t < \infty\}$ on the path space is defined by

$$B_t = \int_0^t b_Q(X_s) ds = \int_0^t \sum_{y \neq X_s} w_{X_s, y} g(X_s, y) ds.$$

Then, the relationship between the functionals A and B is given by the Lévy formula:

$$E_x \sum_{0 < s \leq t} g(X_{s-}, X_s) \alpha(t) = E_x \int_0^t \alpha(t) b_Q(X_s) ds, \quad t > 0,$$

for any continuous positive function $\alpha(t)$.

Now, taking $\alpha(t) = 1$, the Lévy formula tells us

$$E_x A_t = E_x B_t = \int_0^t P(s) \cdot b_Q(x) ds.$$

Furthermore, let

$$g(x, y) = \begin{cases} 1, & \text{if } x \neq y, \\ 0, & \text{if } x = y. \end{cases}$$

Then A_t is the number of transitions of states of X_t up to time t , i. e. $E A_t = N(t)$.

We start at the vertex x . Then

$$E_x A_t = \int_0^t P(s) \cdot b_Q(x) ds = \int_0^t \sum_{y \in V(G)} p_{xy}(s) w_y ds.$$

If we start at an initial distribution θ on graph G , then

$$E_\theta A_t = \int_0^t \theta P(s) \cdot b_Q ds = \int_0^t \sum_{x, y} \theta_x p_{xy}(s) w_y ds.$$

Thus, we have

$$\begin{aligned} \lim_{t \rightarrow \infty} \frac{E_\theta A_t}{t} &= \lim_{t \rightarrow \infty} \frac{\int_0^t \sum_{x,y} \theta_x p_{xy}(s) w_y ds}{t} \\ &= \lim_{t \rightarrow \infty} \sum_{x,y} \theta_x p_{xy}(t) w_y \\ &= \sum_{x,y} \theta_x u_y w_y = \sum_y u_y w_y \end{aligned}$$

and it is independent of the initial condition θ . So,

$$f = \sum_{y \in V(G)} \frac{1}{n} w_y = \frac{2w}{n}.$$

□

Using the notion of frequency, we can compare various Markov times for the continuous time Markov process and its jump chain. Let us recall the notion of Markov time (or stopping time) first. Associated with a stochastic process, there are random variables independent of the future. This kind of random variables are called Markov time or stopping time. Specifically, let σ be a non-negative random variable associated with a given process $\{X_t : 0 \leq t \leq \infty\}$. In the other words, σ associates with each sample function X_t a nonnegative number which we denote by $\sigma(X_t)$. Such a random variable σ is said to be a Markov time relative to the process X_t if it has the following property:

If X_t and Y_t are two sample functions of the process such that $X_\tau = Y_\tau$ for $0 \leq \tau \leq s$ and $\sigma(X_t) < s$, then $\sigma(X_t) = \sigma(Y_t)$.

Now, let's state our main result.

Theorem 2.2. *Let T be any Markov time with finite expectation, i.e. $E(T) < \infty$ associated with the continuous Markov process X_t on a weighted graph G . Let N_T be the number of jumps of X_t during the period $[0, T]$. Then, we have*

$$w_m \leq f \leq w_M$$

and

$$w_m E(T) \leq E(N_T) \leq w_M E(T),$$

where $w_m = \min\{w_x; x \in V(G)\}$ and $w_M = \max\{w_x; x \in V(G)\}$.

Proof. By the Lévy formula, for an initial distribution θ , we have

$$N(t) = E_\theta A_t = \int_0^t \sum_{x,y \in V(G)} \theta_x p_{xy}(s) w_y ds.$$

Since

$$\sum_{x,y \in V(G)} \theta_x p_{xy}(s) w_y \leq \sum_{x,y \in V(G)} \theta_x p_{xy}(s) w_M = w_M$$

and

$$\sum_{x,y \in V(G)} \theta_x p_{xy}(s) w_y \geq \sum_{x,y \in V(G)} \theta_x p_{xy}(s) w_m = w_m,$$

we get

$$w_m t \leq N(t) \leq w_M t.$$

This is

$$w_m \leq \frac{N(t)}{t} \leq w_M.$$

By taking limit, we have

$$w_m \leq f \leq w_M.$$

We suppose that Markov time T has distribution $F(t) = \Pr\{T < t\}$. Then we have

$$\int_0^\infty w_m t dF \leq \int_0^\infty \frac{N(t)}{t} t dF \leq \int_0^\infty w_M t dF.$$

This actually is

$$w_m \int_0^\infty t dF \leq \int_0^\infty N(t) dF \leq w_M \int_0^\infty t dF.$$

We recall the conditional expectation

$$E(N_T) = E(E(A_T | T = t)) = \int_0^\infty N(t) dF.$$

Therefore, we get

$$w_m E(T) \leq E(N_T) \leq w_M E(T).$$

□

The following are two interesting corollaries. The proofs of them are obvious, so we just state the results.

Corollary 2.1. *For CTSRW on a graph G and any Markov time T with finite expectation, we have*

$$d_m \leq f \leq d_M$$

and

$$d_m E(T) \leq E(N_T) \leq d_M E(T).$$

Corollary 2.2. *If G is a regular graph with constant degree d at each vertex, then $f = d$ and $E(N_T) = f E(T)$ for any Markov time T associated with CTSRW on G .*

We may call the inequality in Theorem 2.3 and Corollary 2.1 “time-step inequality”. Of course, we have another version as

$$\frac{E(N_T)}{d_M} \leq E(T) \leq \frac{E(N_T)}{d_m}.$$

In a sense, those inequalities characterize the timing difference between CTSRW and SRW on a graph. It is also interesting to see that the frequency of an unweighted graph is the average of the eigenvalues of its Laplacian. Let $\lambda_1 < \lambda_2 \leq \lambda_3 \leq \dots \leq \lambda_n$ be the spectrum of the Laplacian $L(G)$ of a graph G , $L(G) = D(G) - A(G)$. Then, $\sum_{i=1}^n \lambda_i = \sum_{x \in V(G)} d(x) = 2m$. Thus

$$f = \frac{\sum_{i=1}^n \lambda_i}{n}.$$

Theorem 2.3. *Let G be a non-bipartite graph, and T_x be the first return time of CTSRW on G . Then $E(N_{T_x}) = fE(T_x)$.*

Proof. By the theory of discrete time simple random walks on a graph G , we know $E(N_{T_x}) = \frac{2m}{d(x)}$.

For CTSRW on G , we know $E(T_x) = \frac{n}{d(x)}$, and also $f = \frac{2m}{n}$. Thus

$$E(N_{T_x}) = \frac{2m}{d(x)} = \frac{n}{d(x)} \cdot \frac{2m}{n} = fE(T_x).$$

□

Now, we consider a special problem as that in SRW. Let G be a connected non-bipartite graph. We start our CTSRW at a vertex x and fix a neighboring vertex y of x . What is the expected time that our CTSRW should take in order to return to x through the edge yx ?

For SRW on G , we know the corresponding quantity, the expected number of steps one should take in order to return to x through the edge yx , is $2m$ [Bollobás’].

To deal with the problem for CTSRW on G , we formulate the following Markov time:

$$T_{(x, \overline{yx})} = \inf \{t + \rho_y : t > 0, X(s+t) = y, X(s+t+\rho_y) = x \mid X(s) = x\}.$$

This is the fixed edge first return time, then $E(N_{T_{(x, \overline{yx})}}) = 2m$. By our time-step inequality, we have the following corollary.

Corollary 2.3. *The mean fixed edge first return time has bounds as*

$$\frac{2m}{d_M} \leq E(T_{(x, \overline{yx})}) \leq \frac{2m}{d_m}.$$

The following is another case where the frequency gives us a perfect scaling factor between corresponding quantities of CTSRW and SRW, respectively.

Lemma 2.3. *Let G be a connected graph of order n and size m . The mean hitting time $h(x, y)$ of the CTSRW on G satisfy*

$$\sum_{x \in V(G)} \sum_{y \in \Gamma(x)} h(y, x) = n(n-1).$$

Proof. We know

$$h(x, x) = \frac{1}{d(x)} + \frac{1}{d(x)} \sum_{y \in \Gamma(x)} h(y, x) = \frac{n}{d(x)}.$$

So, we have $\sum_{y \in \Gamma(x)} h(y, x) = n - 1$ which is independent of x . Thus, for n vertices, we will have

$$\sum_{x \in V(G)} \sum_{y \in \Gamma(x)} h(y, x) = n(n - 1).$$

□

If we denote the hitting time of y from x in SRW by $H(x, y)$, then we have an equality

$$\sum_{x \in V(G)} \sum_{y \in \Gamma(x)} H(y, x) = 2m(n - 1).$$

See [2]. So, we have the following theorem.

Theorem 2.4. *With notations as the above, we have*

$$\sum_{x \in V(G)} \sum_{y \in \Gamma(x)} H(y, x) = f \sum_{x \in V(G)} \sum_{y \in \Gamma(x)} h(y, x).$$

Also, for CTSRW on a graph G , we define the mean commute time between vertices x and y to be $c(x, y) = h(x, y) + h(y, x)$. Let $C(x, y)$ be the corresponding quantity for SRW on G . Then we have another version of the above equation in Theorem 2.4 as

$$\sum_{xy \in E(G)} C(x, y) = f \sum_{xy \in E(G)} c(x, y).$$

3. MULTI-PERSON SIMPLE RANDOM WALKS ON GRAPHS

We are led to the multi-person simple random walks (MPSRW) on a graph G by the study of a continuous time Markov process induced by CTSRW on G . The combinatorics of MPSRW is much richer than we have touched upon here.

Let I_n be a finite set of cardinality n . For example, we may have $I_n = \{1, 2, \dots, n\}$. We denote the set of all maps from I_n to itself by M_n . We have the symmetric group S_n sitting inside of M_n . A map $x \in M_n$ is called a *generalized permutation of deficiency k* if

$$|x(I_n)| = n - k.$$

We denote $\text{def}(x) = k$.

M_n is a semigroup under composition. The symmetrical group S_n is a subgroup of M_n . The deficiency determines a grading on M_n which is compatible with the semigroup product on M_n :

$$\text{def}(x) + \text{def}(y) \geq \text{def}(x \circ y).$$

Denote $M_n^{(k)} = \{\text{all maps with deficiency } k\}$, then we have a decomposition of M_n according the deficiency:

$$M_n = M_n^{(0)} \amalg M_n^{(1)} \amalg M_n^{(2)} \amalg \cdots \amalg M_n^{(n-1)},$$

where $M_n^{(0)} = S_n$.

Let G be a graph with the set of vertices $V(G)$ identified with I_n . We will call G the *ground graph*. Let the adjacency matrix $A(G) = (a_{ij})_{n \times n}$ with entries given by

$$a_{ij} = \begin{cases} 1 & \text{if } ij \in E(G), \\ 0 & \text{otherwise.} \end{cases}$$

We define a new graph $M(G)$ as follows: The set of vertices of $M(G)$ is M_n ; For $x, y \in M_n$, there is an edge xy in $M(G)$ only when

$$|\{i : x(i) \neq y(i)\}| = 1,$$

and if $x(i) \neq y(i)$, then $a_{x(i)y(i)} = 1$. We will see that there is a close relationship between the graph $M(G)$ and the n -th tensor power of CTSRW on the ground graph G .

3.1. The tensor product of Markov processes. Let $X_t^{(1)}, X_t^{(2)}, \dots, X_t^{(n)}$ be Markov processes on the state spaces $S^{(1)}, S^{(2)}, \dots, S^{(n)}$ respectively. We define a new process Y_t on the state space $S^{(1)} \times S^{(2)} \times \cdots \times S^{(n)}$ with the transition probability given by

$$\begin{aligned} & \Pr \{Y_{t+h} = (s_2) \mid Y_h = (s_1)\} \\ &= \prod_{k=1}^n \Pr \{Y_{t+h}^{(k)} = s_2^{(k)} \mid Y_h^{(k)} = s_1^{(k)}\} \\ &= \prod_{k=1}^n p_{s_1^{(k)} s_2^{(k)}}^{(k)}(t), \end{aligned}$$

where $(s_i) = (s_i^{(1)}, s_i^{(2)}, \dots, s_i^{(n)})$, $i = 1, 2$, and $p_{s_1^{(k)} s_2^{(k)}}^{(k)}(t)$ is the transition probability of the Markov process $X_t^{(k)}$, $k = 1, 2, \dots, n$.

We call Y_t the *tensor product* of Markov processes $X_t^{(k)}$, $k = 1, 2, \dots, n$.

The next two lemmas can be proved by some direct computations. So we omit the proofs.

Lemma 3.1. Y_t is a Markov process.

Lemma 3.2. *Let $q_{(s_1)(s_2)}$ be the infinitesimal generator of Y_t . Then*

$$q_{(s_1)(s_2)} = \begin{cases} q_{s_1^{(k)} s_2^{(k)}}^{(k)} & \text{if } \exists \text{ only one index } k \text{ such that } s_1^{(k)} \neq s_2^{(k)}, \\ \sum_{k=1}^n q_{s_1^{(k)} s_2^{(k)}}^{(k)} & \text{if } (s_1) = (s_2), \text{ namely, } s_1^{(k)} = s_2^{(k)} \text{ for all } k, \\ 0 & \text{otherwise.} \end{cases}$$

where $q_{s_1^{(k)} s_2^{(k)}}^{(k)}$ is the infinitesimal generator of Markov process $X_t^{(k)}$, $k = 1, 2, \dots, n$.

Now, if we order the elements of the state space $S^{(1)} \times S^{(2)} \times \dots \times S^{(n)}$ lexicographically, it is easy to see that the probability transition matrix of Y_t is given by the tensor product

$$P^{(1)}(t) \otimes P^{(2)}(t) \otimes \dots \otimes P^{(n)}(t).$$

This is why we call Y_t is the tensor product of Markov process $X_t^{(k)}$, $k = 1, 2, \dots, n$. By one of the lemmas above, we also can see that the infinitesimal generator matrix of Y_t is given by

$$Q^{(1)} \otimes I \otimes \dots \otimes I + I \otimes Q^{(2)} \otimes I \otimes \dots \otimes I + \dots + I \otimes \dots \otimes I \otimes Q^{(n)}$$

where $Q^{(k)}$ is the infinitesimal generator matrix of $X_t^{(k)}$, $k = 1, 2, \dots, n$. Thus, for convenience, we denote $Y_t = X_t^{(1)} \otimes X_t^{(2)} \otimes \dots \otimes X_t^{(n)}$.

3.2. CTSRW on $M(G)$. On the graph $M(G)$, we have the Markov process CTSRW. Denote it by Y_t . Let X_t be the Markov process CTSRW on G .

Lemma 3.3. *The Markov process Y_t on the graph $M(G)$ is the n -th tensor power of the Markov process X_t on the ground graph G . The jump chain of Y_t is MPSRW of G .*

Proof. Let's recall $Q = -L(G) = -D(G) + A(G) = (q_{ij})_{n \times n}$,

$$q_{ij} = \begin{cases} 1 & \text{if } ij \in E(G), \\ -d(i) & \text{if } i = j, \\ 0 & \text{otherwise,} \end{cases}$$

where $d(i)$ is the degree of the vertex i . Let $X_t \otimes X_t \otimes \dots \otimes X_t = Z_t$ be the n -th tensor power of the process X_t . Take two states for Z_t , (s_1) and (s_2) . Then (s_i) actually is a sequence of vertices of G . We write $(s_1) = (i_1, i_2, \dots, i_n)$ and $(s_2) = (j_1, j_2, \dots, j_n)$. By the lemma above, we have

$$\begin{aligned} p'_{(s_1)(s_2)}(0) &= \begin{cases} q_{s_1^{(k)} s_2^{(k)}}^{(k)} & \text{if } \exists \text{ only one index } k \text{ such that } s_1^{(k)} \neq s_2^{(k)}, \\ \sum_{k=1}^n q_{s_1^{(k)} s_2^{(k)}}^{(k)} & \text{if } (s_1) = (s_2), \text{ namely, } s_1^{(k)} = s_2^{(k)} \text{ for all } k, \\ 0 & \text{otherwise} \end{cases} \\ &= \begin{cases} 1 & \text{if } \exists \text{ only one index } k \text{ such that } i_k \neq j_k, i_k j_k \in E(G), \\ -\sum_{k=1}^n d(i_k) & \text{if } i_k = j_k \text{ for all } k, \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Now, we can identify (s_1) and (s_2) with the images of certain maps x and y , respectively. Then by the definition of $M(G)$, $xy \in E(M(G))$, if and only if $p'_{(s_1)(s_2)}(0) = 1$. As to the degree of x , we know that the neighbors of $x = (s_1)$ can only be $(k_1, i_2, \dots, i_n)$, where k_1 must be a neighbor of i_1 ; $(i_1, k_2, \dots, i_n)$, where k_2 must be a neighbor of i_2 ; and so on, up to the last one $(i_1, i_2, \dots, i_{n-1}, k_n)$, where k_n must be a neighbor of i_n . Thus

$$d(x) = d(s_1) = \sum_{j=1}^n d(i_j).$$

This agree with $p'_{(s_1)(s_1)}(0) = -\sum_{k=1}^n d(i_k)$. So, $Y_t = X_t \otimes X_t \otimes \dots \otimes X_t = Z_t$.

The second conclusion is easy to see. $\square$

Because of this lemma, we may call graph $M(G)$ the tensor power of the graph G . We also note that the transition probability from x to y in M_n is given by

$$P_{xy}(t) = \prod_{i=1}^n p_{x(i)y(i)}(t).$$

From the proof of Lemma 3.3, we know that for $x \in M_n = V(M(G))$,

$$d(x) = \sum_{k=1}^n d(x(k)).$$

We can compute the size of $M(G)$, i. e. the number of edges of $M(G)$ as

$$\begin{aligned} 2|E(M(G))| &= \sum_{x \in M_n} d(x) = \sum_{x \in M_n} \sum_{k=1}^n d(x(k)) = \sum_{k=1}^n \sum_{x \in M_n} d(x(k)) \\ &= \sum_{k=1}^n ((d(1) + d(2) + \dots + d(n)) n^{n-1}) \\ &= 2mn^n. \end{aligned}$$

Thus, the size of $M(G)$ is mn^n . It is clear that the order of $M(G)$ is n^n . Therefore, the frequency of $M(G)$ is $2m$.

Lemma 3.4. *G is bipartite if and only if $M(G)$ is bipartite.*

Proof. We use the classical result of König that a graph is bipartite if and only if all its cycles are even.

If G is not bipartite, then there is a cycle $C = v_1 v_2 \dots v_m v_1$ of odd length in G . We look at the cycle in $M(G)$ given by

$$(v_1, v_1, \dots, v_1) (v_1, \dots, v_1, v_2) (v_1, \dots, v_1, v_3) \dots (v_1, \dots, v_1, v_m) (v_1, \dots, v_1, v_1).$$

Its length is also odd. Thus $M(G)$ is not bipartite.

If G is bipartite, then the set of vertices $V = V(G)$ can be written as $V_1 \cup V_2$, with $V_1 \cap V_2 = \emptyset$ and there is no edge between vertices both in V_1 or both in V_2 .

We try to bipart the set of vertices of $M(G)$. We know $V(M(G)) = V^{\times n}$. We write $V = V_1 + V_2$. Then

$$V^{\times n} = (V_1 \cup V_2)^{\times n} = V_1^n \cup V_1^{n-1}V_2 \cup V_1^{n-2}V_2^2 \cup \dots \cup V_1V_2^{n-1} \cup V_2^n,$$

where $V_1^{n-k}V_2^k$ means we take $(n-k)$ vertices from V_1 and k vertices from V_2 , regardless of order, to form a vertex of $M(G)$. Let

$$\begin{aligned}\overline{V_1} &= V_1^n \cup V_1^{n-2}V_2^2 \cup V_1^{n-4}V_2^4 \cup \dots, \\ \overline{V_2} &= V_1^{n-1}V_2 \cup V_1^{n-3}V_2^3 \cup V_1^{n-5}V_2^5 \cup \dots.\end{aligned}$$

Then $\overline{V_1} \cup \overline{V_2} = V(M(G))$ and $\overline{V_1} \cap \overline{V_2} = \emptyset$. By the definition of $M(G)$, we can not find an edge between any two vertices which are both in $\overline{V_1}$ or both in $\overline{V_2}$. $\square$

If G is not bipartite with order n and size m . Write the degree sequence of G as $d_1 \leq d_2 \leq \dots \leq d_n$. For any vertex x in $M(G)$, the x -component of the ergodic vector for SRW on $M(G)$ is given by

$$\pi_x = \frac{d(x)}{2mn^n} = \frac{\sum_{i=1}^n d(x(i))}{2mn^n} = n^{-n} \sum_{i=1}^n \frac{d(x(i))}{2m} = n^{-n} \sum_{i=1}^n \pi_{x(i)}^G,$$

where

$$\pi_{x(i)}^G = \frac{d(x(i))}{2m}$$

is the $x(i)$ -component of the ergodic vector for SRW on G .

The expected first return time for CTSRW on $M(G)$ is given by

$$h(x, x) = \frac{n^n}{d(x)} = \frac{n^n}{\sum_{k=1}^n d(x(k))}.$$

In particular, for the identity map Id , $d(Id) = d_1 + d_2 + \dots + d_n = 2m$. So

$$h(Id, Id) = \frac{n^n}{2m}.$$

For the jump chain SRW on $M(G)$, or MPSRW on G , we have

$$H(x, x) = f_{M(G)} h(x, x) = \frac{2mn^n}{\sum_{k=1}^n d(x(k))},$$

where $f_{M(G)} = 2m$ is the frequency of $M(G)$. In particular,

$$H(Id, Id) = n^n.$$

3.3. Hitting time for CTSRW on $M(G)$. For CTSRW on $M(G)$, we are interested in calculating $h(Id, c_i)$, where c_i is the constant map to a vertex i of G . Once we know $h(Id, i)$, we can use it to estimate $H(Id, c_i)$ for SRW on $M(G)$, or equivalently, the expect number of steps MPSRW on G should take to have all persons meet at the vertex i . Namely, we have

$$nd_1 h(Id, c_i) \leq H(Id, c_i) \leq nd_n h(Id, c_i).$$

Notice that when the graph G is regular, $H(Id, c_i)$ is determined completely by $h(Id, c_i)$: $H(Id, c_i) = nd \cdot h(Id, c_i)$.

Let x and y be two distinct vertices in $M(G)$, we consider the hitting time $h(x, y)$ of y from x in CTSRW. We have

$$\begin{aligned} h(x, y) &= n^n \int_0^\infty (P_{yy}(t) - P_{xy}(t)) dt \\ &= n^n \int_0^\infty \left(\prod_{i=1}^n p_{y(i)y(i)}(t) - \prod_{i=1}^n p_{x(i)y(i)}(t) \right) dt. \end{aligned}$$

For $x = Id$, $y = c_i$, we have

$$h(Id, c_i) = n^n \int_0^\infty \left(p_{ii}(t)^n - \prod_{j=1}^n p_{ji}(t) \right) dt.$$

To calculate this integral, we diagonalize the Laplacian of G . Write

$$U^T Q U = \text{diag}[-\lambda_1, -\lambda_2, \dots, -\lambda_n],$$

where $U = (u_{ij})_{n \times n}$ is an orthogonal matrix. Then

$$p_{ij}(t) = \sum_{k=1}^n u_{ik} u_{jk} e^{-\lambda_k t}.$$

Since

$$p_{ii}(t)^n = \left(\sum_{k=1}^n u_{ik}^2 e^{-\lambda_k t} \right)^n = \sum_{1 \leq k_1, k_2, \dots, k_n \leq n} u_{ik_1}^2 u_{ik_2}^2 \cdots u_{ik_n}^2 e^{-(\lambda_{k_1} + \lambda_{k_2} + \cdots + \lambda_{k_n})t}$$

and

$$\prod_{k=1}^n p_{ki}(t) = \sum_{1 \leq k_1, k_2, \dots, k_n \leq n} u_{1k_1} u_{ik_1} u_{2k_2} u_{ik_2} \cdots u_{nk_n} u_{ik_n} e^{-(\lambda_{k_1} + \lambda_{k_2} + \cdots + \lambda_{k_n})t},$$

we get

$$h(Id, c_i) = n^n \sum_{1 \leq k_1, k_2, \dots, k_n \leq n} \frac{u_{ik_1}^2 u_{ik_2}^2 \cdots u_{ik_n}^2 - u_{1k_1} u_{ik_1} u_{2k_2} u_{ik_2} \cdots u_{nk_n} u_{ik_n}}{\lambda_{k_1} + \lambda_{k_2} + \cdots + \lambda_{k_n}}.$$

Example. Let G be the triangle graph. It is regular with $n = 3$ and $d = 2$. The matrix Q and U are given below:

$$Q = \begin{pmatrix} -2 & 1 & 1 \\ 1 & -2 & 1 \\ 1 & 1 & -2 \end{pmatrix} \quad \text{and} \quad U = \begin{pmatrix} -\frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} \\ 0 & \frac{2}{\sqrt{6}} & \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} \end{pmatrix}.$$

We have $U^T Q U = \text{diag}[-3, -3, 0]$. Since $\lambda_3 = 0$, when calculating $h(Id, c_i)$, we should drop the term $(k_1, k_2, k_3) = (3, 3, 3)$. Notice that the numerator of this term in $h(Id, c_i)$ is also zero, so it is fine to drop this term. We have

$$h(Id, c_1) = 3^3 \cdot \frac{31}{162} = \frac{31}{6}.$$

So

$$H(Id, c_1) = 3 \cdot 2 \cdot \frac{31}{6} = 31.$$

Thus, for our MPSRW on the triangle graph, it takes 31 steps on average for 3 persons to meet at any specified vertex, given that they all start at different vertices.

For the square graph with $n = 4$ and $d = 2$, a similar calculation shows

$$h(Id, c_1) = 4^4 \cdot \frac{167}{1120} = \frac{1336}{35}$$

and

$$H(Id, c_1) = 8 \cdot \frac{1336}{35} \approx 305.371.$$

Thus, for our MPSRW on the square graph, it takes about 305 steps on average for 4 persons to meet at any specified vertex, given that they all start at different vertices.

REFERENCES

- [1] D. Aldous, *Hitting times for random walks on vertex-transitive graphs*, Math. Proc. Camb. Phil. Soc. **106**(1989), pp. 179–191.
- [2] B. Bollobás, *Modern Graph Theory*, Springer, New York, 1998.
- [3] F. Buckley and F. Harary, *Distance in Graphs*, Addison-Wesley, 1990.
- [4] K.-L. Chung, *Markov Chains with Stationary Transition Probabilities*, Springer-Verlag, Berlin, 1967.
- [5] C. Godsil and G. Royle, *Algebraic Graph Theory*, Springer, New York, 2001.
- [6] L. Lovász, *Random walks on graphs: a survey*, Bolyai Society Mathematical Study, Combinatorics, **2**(1993), pp. 1–46.
- [7] R. Syski, *Passage Times for Markov Chains*, IOS Press, Amsterdam, 1992.

MATHEMATICAL BIOSCIENCES INSTITUTE, THE OHIO STATE UNIVERSITY, COLUMBUS, OH
43210, USA

E-mail address: `tianjj@mbi.ohio-state.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, RIVERSIDE, CA 92521,
USA

E-mail address: `xl@math.ucr.edu`